

wherein, the ring A represents an aryl ring,

R<sup>1</sup> represents a hydroxy group, an amino group, or a lower alkylamino group having 1 to 4 carbons that may be substituted with a carboxylic group, a lower aralkylamino group having 7 to 10 carbons that may be substituted with a carboxylic group, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, a lower alkyl group having 1 to 4 carbons substituted with a carboxylic group, or a lower alkylene group having 2 to 4 carbons substituted with a carboxylic group;

R<sup>2</sup> and R<sup>3</sup>, which may be the same or different, represent a hydrogen, a lower alkyl group having 1 to 4 carbons that may be substituted, a halogen atom, a hydroxy group, a

contd.  
B2


~~lower alkoxy group having 1 to 4 carbons, an amino group, a lower alkylamino group having 1 to 4 carbons that may be substituted, a lower aralkylamino group having 7 to 10 carbons that may be substituted, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, or a carboxylic group; or~~

~~when the ring A is a benzene ring, R<sup>1</sup> and R<sup>2</sup>, together with the benzene ring to be substituted, may form a fused heterocyclic ring that may be substituted with a carboxylic acid, and a carbon atom in said fused heterocyclic ring may form a carbonyl group wherein R<sup>3</sup> is as defined above; and~~

~~X represents a hydrogen atom, a lower alkyl group having 1 to 4 carbons, a lower alkoxy group having 1 to 4 carbons, a halogen atom, a hydroxy group, an amino group, or a nitro group;~~

~~or a pharmaceutically acceptable salt thereof.~~

7. (Amended) The pharmaceutical composition according to claim 3, wherein said chymase inhibitor is a quinazoline derivative represented by the formula (I):



wherein, the ring A represents an aryl ring,


~~R<sup>1</sup> represents a hydroxy group, an amino group, or a lower alkylamino group having 1 to 4 carbons that may be substituted with a carboxylic group, a lower aralkylamino group having 7 to 10 carbons that may be substituted with a carboxylic group, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, a lower alkyl group having 1 to 4 carbons substituted with a carboxylic group, or a lower alkylene group having 2 to 4 carbons substituted with a carboxylic group;~~

~~SECRET~~

X represents a hydrogen atom, a lower alkyl group having 1-4 carbons, a lower alkoxy group having 1 to 4 carbons, a halogen atom, a hydroxy group, an amino group, or a nitro group;  
or a pharmaceutically acceptable salt thereof.

**Please add new claims 9-10 as follows.**

--9. The preventive or therapeutic agent according to claim 2, wherein said chymase inhibitor is a quinazoline derivative represented by the formula (1):



wherein, the ring A represents an aryl ring,

R<sup>1</sup> represents a hydroxy group, an amino group, or a lower alkylamino group having 1 to 4 carbons that may be substituted with a carboxylic group, a lower aralkylamino group having 7 to 10 carbons that may be substituted with a carboxylic group, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, a lower alkyl group having 1 to 4 carbons substituted with a carboxylic group, or a lower alkylene group having 2 to 4 carbons substituted with a carboxylic group;


that may  
aromat.

$\mathbf{R}^3$  is as defined above; and

a nitro group;

or a pharmaceutically acceptable salt thereof.

10. The pharmaceutical composition according to claim 4, wherein said chymase inhibitor is a quinazoline derivative represented by the formula (1):



wherein, the ring A represents an aryl ring,

*AD*  
R<sup>1</sup> represents a hydroxy group, an amino group, or a lower alkylamino group having 1 to 4 carbons that may be substituted with a carboxylic group, a lower aralkylamino group having 7 to 10 carbons that may be substituted with a carboxylic group, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, a lower alkyl

group having 1 to 4 carbons substituted with a carboxylic group, or a lower alkylene group having 2 to 4 carbons substituted with a carboxylic group;

$R^2$  and  $R^3$ , which may be the same or different, represent a hydrogen, a lower alkyl group having 1 to 4 carbons that may be substituted, a halogen atom, a hydroxy group, a lower alkoxy group having 1 to 4 carbons, an amino group, a lower alkylamino group having 1 to 4 carbons that may be substituted, a lower aralkylamino group having 7 to 10 carbons that may be substituted, an amino group acylated with a lower fatty acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group acylated with an aromatic carboxylic acid that may be substituted with a carboxylic group, an amino group acylated with a heteroaromatic carboxylic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a lower alkanesulfonic acid having 1 to 4 carbons that may be substituted with a carboxylic group, an amino group sulfonylated with an aromatic sulfonic acid that may be substituted with a carboxylic group, an amino group sulfonylated with a heteroaromatic sulfonic acid that may be substituted with a carboxylic group, or a carboxylic group; or

when the ring A is a benzene ring,  $R^1$  and  $R^2$ , together with the benzene ring to be substituted, may form a fused heterocyclic ring that may be substituted with a carboxylic acid, and a carbon atom in said fused heterocyclic ring may form a carbonyl group wherein  $R^3$  is as defined above; and

X represents a hydrogen atom, a lower alkyl group having 1-4 carbons, a lower alkoxy group having 1 to 4 carbons, a halogen atom, a hydroxy group, an amino group, or a nitro group;